Total Marks: 100

5

6

8

6

8

6

N.B.: (1) Question No. 1 is compulsory.

(2) Attempt any four questions from question Nos. 2 to 7.

(3) If in doubt make suitable assumption. Justify your assumption and proceed.

Significant full marks.

(4) Figures to the right indicate full marks.

(a) If the angle between the surfaces
$$x^2 + axz + byz = 2$$
 and $x^2z + xy + y + 1 = z$ at

(0, 1, 2) is $\cos^{-1} \frac{1}{\sqrt{2}}$ then find the constants a and b.

(b) Find the eigen values and eigen vectors of the orthogonal matrix :-

 $B = \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \end{bmatrix}$

(c) Evaluate
$$\int_{c}^{c} f(z) dz$$
 along the parabola $y = 2x^2$ from $z = 0$ to $z = 3 + 18i$ where 5 $f(z) = x^2 - 2ixy$.

Find unit normal vector to the unit sphere at point -

 $\left(\begin{array}{c} a & a & a \\ \sqrt{3} & \sqrt{3} & \sqrt{3} \end{array}\right)$

2. (a) Find the directional derivative of
$$xy^2 + yz^3$$
 at the point (2, -1, 1) along the tangent to 6 the curve $x = a \sin t$, $y = a \cos t$, $z = at$ at $t = \frac{\pi}{4}$.

(b) Verify Cauchy's integral theorem for $f(z) = e^z$ along a circle c : |z| = 1.

(c) Reduce the Quadratic form - $8x^2 + 7y^2 + 3z^2 + 12xy + 4xz - 8yz$ to sum of squares and find the corresponding Linear transformation also find the rank, index and signature.

Using Caley-Hamilton theorem for -3. (a)

 $A = \begin{bmatrix} -3 & -4 & 4 \\ 2 & 2 & 1 \\ 0 & 1 & 2 \end{bmatrix}$

Find A64 + 2A37 - 581.

(b) Prove that
$$\nabla^2 f(r) = f''(r) + \frac{2}{r}f'(r)$$
 and hence show that $\nabla^4 e^r = \left(1 + \frac{4}{r}\right)e^r$.

Find all possible Laurent's expansion of the function :-

 $f(z) = \frac{7z-2}{z(z-2)(z+1)}$ about z = -1.

4. (a) If
$$A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$
 and $B = \begin{bmatrix} 2 & 0 \\ 1/2 & 2 \end{bmatrix}$ then prove that both A and B are not diagonalizable 6 but AB is not diagonalizable.

(b) Verify Green's theorem in plane for :-

 $\oint (x^2 - 2xy) dx + (x^2y + 3) dy$ where c is the boundary of the region defined

http://www.howtoexam.com

Con. 3436-CO-9712-08. Sum IV Apr Math IV 1916/08

(c) (i) Evaluate
$$\int_{-\infty}^{\infty} \frac{x^2}{(x^2+a^2)(x^2+b^2)} dx$$
, $a>0$, $b>0$

(c) (i) Evaluate
$$\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + a^2)(x^2 + b^2)} dx$$
, $a > 0$, $b > 0$

(ii) Evaluate
$$\int_{0}^{2\pi} \frac{d\theta}{\sqrt{2-\cos\theta}}$$

5. (a) Evaluate
$$\int_{c} \frac{\sin^{6} z}{(z - \pi/6)^{3}} dz \text{ where c is } |z| = 1.$$

$$A = \begin{bmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{bmatrix}$$
 is derogatory.

(c) Verify Gauss-Divergence theorem for :-
$$F = 4xi + 2y^2j + z^2k \text{ taken over the region of the cylinder bounded by } x^2 : y^2 = 4,$$

$$z = 0 \text{ and } z = 3.$$

6. (a) If
$$A = \begin{bmatrix} -1 & 4 \\ 2 & 1 \end{bmatrix}$$
 then prove that 3 tan $A = A$ tan 3.

(b) Evaluate
$$\int_{c} (z-z^2) dz$$
 where c is the upper half of circle $|z-2|=3$.

(c) (i) Show that
$$\vec{F} = (ye^{xy}cosz)i + (xe^{xy}cosz)j + (-e^{xy}sinz)k$$
 is irrotational and 6 find the scalar potential ϕ such that $\vec{F} = \nabla \phi$

(ii) Find div F where
$$\overrightarrow{F} = \frac{xi - yj}{x^2 + y^2}$$

7. (a) State and prove Cauchys-Residue theorem and hence – 6

Evaluate
$$\int \frac{1+z}{z(2-z)} dz$$
 where c is $|z| = 1$.

(b) Evaluate
$$\iint_S F \cdot nds$$
 where $F = (x + y^2) i - 2xj + 2yzk$ and s is the surface of the plane 6 $2x + y + z = 6$ in the first octant.

$$A = \begin{bmatrix} -9 & 4 & 4 \\ -8 & 3 & 4 \\ -16 & 8 & 7 \end{bmatrix}$$