

Name :

Roll No. :

Invigilator's Signature :

**CS/B.Tech(NEW)/SEM-1/M-101/2010-11
2010-11**

MATHEMATICS - I

Time Allotted : 3 Hours

Full Marks : 70

The figures in the margin indicate full marks.

Candidates are required to give their answers in their own words
as far as practicable.

GROUP - A

(Multiple Choice Type Questions)

1. Choose the correct alternatives for any ten of the following :
 $10 \times 1 = 10$

i) If α, β are the roots of the equation $x^2 - 3x + 2 = 0$

then
$$\begin{bmatrix} 0 & \alpha & \beta \\ \beta & 0 & 0 \\ 1 & -\alpha & \alpha \end{bmatrix}$$
 is

a) 6

b) $\frac{3}{2}$

c) - 6

d) 3.

ii) If $y = e^{ax+b}$, then $(y_5)_0 =$

a) ae^b

b) a^5e^b

c) $a^b e^{ax}$

d) none of these.

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

iii) If Rolle's theorem is applied to $f(x) = x(x^2 - 1)$ in $[0, 1]$, then $C =$

- a) 1 b) 0
 c) $-\frac{1}{\sqrt{3}}$ d) $\frac{1}{\sqrt{3}}$.

iv) If $u + v = x$, $uv = y$, then $\frac{\partial(u, v)}{\partial(x, y)} =$

- a) $u - v$ b) uv
 c) $u + v$ d) $\frac{u}{v}$.

v) The value of $\int_{-\pi/2}^{\pi/2} \sin^7 \theta d\theta$ is

- a) $\frac{6.4.2}{7.5.3.1}$ b) $\frac{6}{7}$
 c) 0 d) $\frac{2.(6.4.2)}{7.5.3.1}$.

vi) The sequence $\left\{ (-1)^n \frac{1}{n} \right\}$ is

- a) convergent b) oscillatory
 c) divergent d) none of these.

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

vii) If $\vec{\alpha} = 3\hat{i} - 2\hat{j} + \hat{k}$, $\vec{\beta} = 2\hat{i} - \hat{k}$, then $(\vec{\alpha} \times \vec{\beta}) \cdot \vec{\alpha}$ is equal to

- a) $\hat{i} + \hat{j} + \hat{k}$ b) $\hat{i} + \hat{k}$
c) $\hat{i} - \hat{k}$ d) 0.

viii) The matrix $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is

- a) symmetric b) skew-symmetric
c) singular d) orthogonal.

ix) The value of t for which

$\vec{J} = (x + 3y)\hat{i} + (y - 2z)\hat{j} + (x + tz)\hat{k}$ is solenoidal is

- a) 2 b) -2
c) 0 d) 1.

x) The distance between two parallel planes $x + 2y - z = 4$ and $2x + 4y - 2z = 3$ is

- a) $\frac{5}{\sqrt{24}}$ b) $\frac{5}{24}$
c) $\frac{11}{\sqrt{24}}$ d) none of these.

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

- xi) In the M.V. theorem $f(h) = f(o) + hf'(oh)$; $0 < \theta < 1$ if $f(x) = \frac{1}{1+x}$ and $h = 3$, then value of θ is

- a) 1 b) $\frac{1}{3}$
c) $\frac{1}{\sqrt{2}}$ d) none of these.

- xii) The series $\sum \frac{1}{np}$ is convergent if

- a) $p \geq 1$ b) $p > 1$
c) $p < 1$ d) $p \leq 1$.

GROUP - B

(Short Answer Type Questions)

Answer any three of the following. $3 \times 5 = 15$

2. If $y = (x^2 - 1)^n$, then show that

$$(x^2 - 1)y_{n+2} + 2xy_{n+1} - n(n+1)y_n = 0. \text{ Hence}$$

find $y_n(0)$.

3. Using M.V.T. prove that

$$x > \tan^{-1} x > \frac{x}{1+x^2}, \quad 0 < x < \pi/2.$$

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

4. Show that

$$\begin{bmatrix} 1+a & 1 & 1 & 1 \\ 1 & 1+b & 1 & 1 \\ 1 & 1 & 1+c & 1 \\ 1 & 1 & 1 & 1+d \end{bmatrix} = abcd \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} \right).$$

5. Test the nature of the series

$$\left(\frac{1}{3} \right)^2 + \left(\frac{1 \cdot 2}{3 \cdot 5} \right)^2 + \left(\frac{1 \cdot 2 \cdot 3}{3 \cdot 5 \cdot 7} \right)^2 + \dots \dots$$

6. If \vec{a} , \vec{b} , \vec{c} are three vectors, then show that

$$[\vec{a} \times \vec{b} \quad \vec{b} \times \vec{c} \quad \vec{c} \times \vec{a}] = [a \ b \ c]^2.$$

7. If $u = \tan^{-1} \frac{x^2 - y^2}{x - y}$, show that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{1}{2} \sin 2u$.

GROUP - C

(Long Answer Type Questions)

Answer any three of the following. $3 \times 15 = 45$

8. a) Determine the conditions under which the system of equations $x + y + z = 1$, $x + 2y - z = b$, $5x + 7y + az = b^2$, admits of

- i) only one solution
- ii) no solution
- iii) many solutions.

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

- b) Find the eigenvalues and the corresponding eigen-

vecgors of the matrix $A = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 0 & 0 & 1 \end{pmatrix}$.

- c) Find whether the following series is convergent :

$$\left(\frac{2^2}{1^2} - \frac{2}{1}\right)^{-1} + \left(\frac{3^3}{2^3} - \frac{3}{2}\right)^{-2} + \left(\frac{4^4}{3^4} - \frac{4}{3}\right)^{-3} + \dots$$

9. a) If $f(x) = x^2$, $g(x) = x^3$ on $[1, 2]$, is Cauchy's mean value theorem applicable ? If so, find ξ .

- b) If $I_n = \int \frac{\cos n\theta}{\cos \theta} d\theta$, show that

$$(n-1)(I_n + I_{n-2}) = 2 \sin(n-1)\theta.$$

Hence evaluate $\int (4 \cos^2 \theta - 3) d\theta$.

- c) If $r = |\vec{r}|$, where $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, prove that

$$\nabla(r^n) = nr^{n-2} \vec{r}.$$

10. a) Find $\partial(u, v)$, where $u = x^2 - 2y^2$, $v = 2x^2 - y^2$

$$\partial(r, \theta)$$

and $x = r \cos \theta$, $y = \sin \theta$.

CS/B.Tech(NEW)/SEM-1/M-101/2010-11

b) Verify Green's theorem for $\vec{F} = (xy + y^2)\hat{i} + x^2\hat{j}$

where the curve C is bounded by $y = x$ and $y = x^2$.

c) Evaluate : $\int_0^a \int_0^x \int_0^y x^3 y^2 z \, dz \, dy \, dx.$

11. a) Find the maxima and minima of the function $x^3 + y^3 - 3x + 12y + 20$. Also find the saddle point.

b) State Cayley- Hamilton theorem and verify the same for the matrix $A = \begin{bmatrix} 1 & 2 \\ -2 & -1 \end{bmatrix}$. Find A^{-1} and A^8 .

c) Show that $\text{Curl } \nabla f = 0$,

where $f(x, y, z) = x^2y + 2xy + z^2$.

12. a) Given the function $= \begin{cases} \frac{xy(x^2 - y^2)}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0 & , (x, y) = (0, 0) \end{cases}$

Find from definition $f_{xy}(0, 0)$ and $f_{yx}(0, 0)$.

Is $f_{xy} = f_{yx}$?

b) Integrate by changing the order of integration

$$\int_0^a \int_{x^2/a}^{2a-x} xy \, dy \, dx.$$

c) If $F(p, v, t) = 0$, show that

$$\left(\frac{dp}{dt}\right)_{v \text{ constant}} \times \left(\frac{dv}{dp}\right)_{t \text{ constant}} \times \left(\frac{dt}{dv}\right)_{p \text{ constant}} = -1.$$